

B UK Patent Application (19) GB (11) 2 168 973 A

(43) Application published 2 Jul 1986

(21) Application No 8529799

(22) Date of filing 3 Dec 1985

(30) Priority data

(31) 8430624	(32) 4 Dec 1984	(33) GB
8430625	4 Dec 1984	
8430626	4 Dec 1984	
8430773	5 Dec 1984	

(51) INT CL⁴
C07D 209/14 A61K 31/40

(52) Domestic classification (Edition H):
C2C 1175 1343 200 213 220 225 226 22Y 246 247 250
251 25Y 29X 29Y 304 30Y 323 32Y 342 34Y 364 36Y 385
510 512 51X 537 538 620 650 662 682 802 80Y AA SJ
U1S 2415 C2C

(56) Documents cited
None

(58) Field of search
C2C


(71) Applicant
Glaxo Group Limited (United Kingdom),
Clarges House, 6-12 Clarges Street, London W1Y 8DH

(72) Inventors
Brian Evans,
Alexander William Oxford,
Darko Butina

(74) Agent and/or Address for Service
Elkington and Fife, High Holborn House, 52-54 High
Holborn, London WC1V 6SH

(54) Indole derivatives

(57) Indole derivatives are disclosed of formula (I):



(I)

wherein

R₁ represents H, alkyl or alkenyl;

R₂ represents H, alkyl, alkenyl, cycloalkyl or phenyl or phenyl alkyl the phenyl ring being optionally substituted by halogen, alkyl, alkoxy, hydroxyl or by a group -NR_aR_b, or -CONR_aR_b, wherein R_a and R_b are H, alkyl, alkenyl, or with the nitrogen atom form a saturated monocyclic ring;

R₃ and R₄ are H, alkyl or propenyl or together form an aralkylidene group;

Alk represents a C₂₋₃ alkyl chain optionally substituted by one or two alkyl groups; and

A' represents a C₂₋₆ alkyl or alkenyl chain and salts and solvates thereof.

The compounds have selective vasoconstrictor activity and are useful in treating and/or preventing pain resulting from dilation of the cranial vasculature, particularly migraine.

GB 2 168 973

698
DE 3542 072
24.1.2.4

SPECIFICATION

Indole derivatives


5 This invention relates to indole derivatives, to processes for their preparation, to pharmaceutical compositions containing them and to their medical use, in particular to compounds and compositions of use in the treatment of migraine.

The pain of migraine is associated with excessive dilatation of the cranial vasculature and known treatments for migraine include the administration of compounds having vasoconstrictor properties such as 10 ergotamine. However, ergotamine is a non-selective vasoconstrictor which constricts blood vessels throughout the body and has undesirable and potentially dangerous side effects. Migraine may also be treated by administering an analgesic, usually in combination with an antiemetic, but such treatments are of limited value.

There is thus a need for a safe and effective drug for the treatment of migraine, which can be used 15 either prophylactically or to alleviate an established headache, and a compound having a selective vasoconstrictor activity would fulfil such a role.

We have now found a group of indole derivatives having potent and selective vasoconstrictor activity.

The present invention provides indoles of the general formula (I):



(I)

25 wherein

R represents a hydrogen atom or a C₁-alkyl or C₁-alkenyl group;

R₁ represents a hydrogen atom, a C₁-alkyl, C₁-alkenyl, or C₁-cycloalkyl group, or a phenyl or phenyl 30 (C₁-alkyl group in which the phenyl ring may be unsubstituted or substituted by a halogen atom, a C₁-alkyl, C₁-alkoxy or hydroxyl group, or by a group -NH-R₃, or -CONR₃, wherein R₃ and R₄, which may be the same or different, each represents a hydrogen atom or a C₁-alkyl or C₁-alkenyl group, or together with the nitrogen atom to which they are attached form a saturated monocyclic 5 to 7-membered ring, which may contain an additional hetero function, for example, an oxygen atom or the group NR₅ (where R₅ is a hydrogen atom or a lower alkyl group); R₂ and R₃, which may be the same or different, each 35 represents a hydrogen atom or a C₁-alkyl or propenyl group or R₂ and R₃ together form an aralkylidene group;


Alk represents an alkyl chain containing two or three carbon atoms which may be unsubstituted or substituted by not more than two C₁-alkyl groups; and

A' represents an alkenyl chain containing two to five carbon atoms, and salts and solvates thereof.

40 All optical isomers of compounds of general formula (I) and their mixtures, including the racemic mixtures thereof, are embraced by the invention. The invention also includes within its scope geometric isomers of compounds (I) and mixtures of such isomers.

Referring to the general formula (I), the alkyl groups and the alkyl moiety of the alkoxy groups may be straight chain or branched chain alkyl groups containing 1 to 3 carbon atoms, or in the case of R₁, 1 to 6, 45 preferably 1 to 3, carbon atoms. Examples of alkyl groups include methyl, ethyl, propyl and isopropyl groups. The alkenyl groups preferably contain 3 or 4 carbon atoms, examples of which include propenyl and butenyl groups. The cycloalkyl groups preferably contain 5 or 6 carbon atoms and examples include cyclopentyl and cyclohexyl groups. The alkyl moieties of the phenylalkyl groups preferably contain 1 or 2 carbon atoms as, e.g., benzyl and phenylethyl groups. The aralkylidene group is preferably an aryl 50 methylidene group such as benzylidene. When R₁ represents a substituted phenyl or phenyl (C₁-alkyl group the substituent may be in the ortho, meta or para positions. A halogen substituent on a phenyl ring in general formula (I) may be for example a fluorine, chlorine or bromine atom.


The alkenyl chain A' may for example, be represented by the formula




wherein m is zero or an integer from 1 to 3 and n is zero or an integer from 1 to 3, such that the sum of m and n together does not exceed 3.

When R₁ represents a substituted phenyl or phenyl (C₁-alkyl group, m and n preferably each represent 60 zero, 1 or 2, such that the sum of m and n together does not exceed 2.

It will be appreciated that the compounds of formula (I) may exist in the E- or Z- configuration with respect to the double bond in the alkenyl chain $-(\text{CH}_2)_m-\text{CH}=\text{CH}(\text{CH}_2)_n-$. The present invention includes within its scope both isomeric forms as well as mixtures thereof. In general, compounds of the invention in the E-configuration are preferred. The E-configuration may be represented structurally as:




5 In the compounds of general formula (I), the alkenyl chain A¹ is preferably a group of formula :



wherein m is as previously defined, preferably zero or 1 and n is zero or 1, most preferably zero.

10 Thus, a preferred class of compounds according to the invention is that represented by general formula (I'):



(wherein R₁, R₂, R₃, R₄, Alk and m are as previously defined) and physiologically acceptable salts and solvates (e.g. hydrates) thereof.


20 In the compounds of general formulae (I) and (I') Alk preferably represents an unsubstituted alkyl chain, especially an unsubstituted alkyl chain containing two carbon atoms.

R₁ is preferably a hydrogen atom or a C₁₋₆ alkyl group and R₂ preferably represents a hydrogen atom, a C₁₋₆ alkyl group, a C₃₋₆ cycloalkyl group or a substituted or unsubstituted phenyl or phenyl(C₁₋₆) alkyl group. It is particularly preferred that one of R₁ or R₂ represents a hydrogen atom. When R₁ represents a substituted phenyl or phenyl (C₁₋₆) alkyl group it is preferred that R₁ represents a hydrogen atom or a C₁₋₆ alkyl group.

Preferred substituents on the phenyl or phenyl (C_{10}) alkyl group represented by R_1 are C_{1-4} alkoxy groups and groups of the formula $-CONR_2R_3$ wherein R_2 and R_3 , which may be the same or different each represents a hydrogen atom or a C_{1-4} alkyl group.

30 R_1 and R_2 , which may be the same or different preferably each represent a hydrogen atom or a C_{1-6} alkyl group.

A particularly preferred class of compounds according to the invention is that represented by the general formula (Ia):



wherein

R₁ represents a hydrogen atom or a C₁₋₄ alkyl group (e.g. methyl);

R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, R₂₄, R₂₅, R₂₆, R₂₇, R₂₈, R₂₉, R₃₀, R₃₁, R₃₂, R₃₃, R₃₄, R₃₅, R₃₆, R₃₇, R₃₈, R₃₉, R₄₀, R₄₁, R₄₂, R₄₃, R₄₄, R₄₅, R₄₆, R₄₇, R₄₈, R₄₉, R₅₀, R₅₁, R₅₂, R₅₃, R₅₄, R₅₅, R₅₆, R₅₇, R₅₈, R₅₉, R₆₀, R₆₁, R₆₂, R₆₃, R₆₄, R₆₅, R₆₆, R₆₇, R₆₈, R₆₉, R₇₀, R₇₁, R₇₂, R₇₃, R₇₄, R₇₅, R₇₆, R₇₇, R₇₈, R₇₉, R₈₀, R₈₁, R₈₂, R₈₃, R₈₄, R₈₅, R₈₆, R₈₇, R₈₈, R₈₉, R₉₀, R₉₁, R₉₂, R₉₃, R₉₄, R₉₅, R₉₆, R₉₇, R₉₈, R₉₉, R₁₀₀, R₁₀₁, R₁₀₂, R₁₀₃, R₁₀₄, R₁₀₅, R₁₀₆, R₁₀₇, R₁₀₈, R₁₀₉, R₁₁₀, R₁₁₁, R₁₁₂, R₁₁₃, R₁₁₄, R₁₁₅, R₁₁₆, R₁₁₇, R₁₁₈, R₁₁₉, R₁₂₀, R₁₂₁, R₁₂₂, R₁₂₃, R₁₂₄, R₁₂₅, R₁₂₆, R₁₂₇, R₁₂₈, R₁₂₉, R₁₃₀, R₁₃₁, R₁₃₂, R₁₃₃, R₁₃₄, R₁₃₅, R₁₃₆, R₁₃₇, R₁₃₈, R₁₃₉, R₁₄₀, R₁₄₁, R₁₄₂, R₁₄₃, R₁₄₄, R₁₄₅, R₁₄₆, R₁₄₇, R₁₄₈, R₁₄₉, R₁₅₀, R₁₅₁, R₁₅₂, R₁₅₃, R₁₅₄, R₁₅₅, R₁₅₆, R₁₅₇, R₁₅₈, R₁₅₉, R₁₆₀, R₁₆₁, R₁₆₂, R₁₆₃, R₁₆₄, R₁₆₅, R₁₆₆, R₁₆₇, R₁₆₈, R₁₆₉, R₁₇₀, R₁₇₁, R₁₇₂, R₁₇₃, R₁₇₄, R₁₇₅, R₁₇₆, R₁₇₇, R₁₇₈, R₁₇₉, R₁₈₀, R₁₈₁, R₁₈₂, R₁₈₃, R₁₈₄, R₁₈₅, R₁₈₆, R₁₈₇, R₁₈₈, R₁₈₉, R₁₉₀, R₁₉₁, R₁₉₂, R₁₉₃, R₁₉₄, R₁₉₅, R₁₉₆, R₁₉₇, R₁₉₈, R₁₉₉, R₂₀₀, R₂₀₁, R₂₀₂, R₂₀₃, R₂₀₄, R₂₀₅, R₂₀₆, R₂₀₇, R₂₀₈, R₂₀₉, R₂₁₀, R₂₁₁, R₂₁₂, R₂₁₃, R₂₁₄, R₂₁₅, R₂₁₆, R₂₁₇, R₂₁₈, R₂₁₉, R₂₂₀, R₂₂₁, R₂₂₂, R₂₂₃, R₂₂₄, R₂₂₅, R₂₂₆, R₂₂₇, R₂₂₈, R₂₂₉, R₂₃₀, R₂₃₁, R₂₃₂, R₂₃₃, R₂₃₄, R₂₃₅, R₂₃₆, R₂₃₇, R₂₃₈, R₂₃₉, R₂₄₀, R₂₄₁, R₂₄₂, R₂₄₃, R₂₄₄, R₂₄₅, R₂₄₆, R₂₄₇, R₂₄₈, R₂₄₉, R₂₅₀, R₂₅₁, R₂₅₂, R₂₅₃, R₂₅₄, R₂₅₅, R₂₅₆, R₂₅₇, R₂₅₈, R₂₅₉, R₂₆₀, R₂₆₁, R₂₆₂, R₂₆₃, R₂₆₄, R₂₆₅, R₂₆₆, R₂₆₇, R₂₆₈, R₂₆₉, R₂₇₀, R₂₇₁, R₂₇₂, R₂₇₃, R₂₇₄, R₂₇₅, R₂₇₆, R₂₇₇, R₂₇₈, R₂₇₉, R₂₈₀, R₂₈₁, R₂₈₂, R₂₈₃, R₂₈₄, R₂₈₅, R₂₈₆, R₂₈₇, R₂₈₈, R₂₈₉, R₂₉₀, R₂₉₁, R₂₉₂, R₂₉₃, R₂₉₄, R₂₉₅, R₂₉₆, R₂₉₇, R₂₉₈, R₂₉₉, R₃₀₀, R₃₀₁, R₃₀₂, R₃₀₃, R₃₀₄, R₃₀₅, R₃₀₆, R₃₀₇, R₃₀₈, R₃₀₉, R₃₁₀, R₃₁₁, R₃₁₂, R₃₁₃, R₃₁₄, R₃₁₅, R₃₁₆, R₃₁₇, R₃₁₈, R₃₁₉, R₃₂₀, R₃₂₁, R₃₂₂, R₃₂₃, R₃₂₄, R₃₂₅, R₃₂₆, R₃₂₇, R₃₂₈, R₃₂₉, R₃₃₀, R₃₃₁, R₃₃₂, R₃₃₃, R₃₃₄, R₃₃₅, R₃₃₆, R₃₃₇, R₃₃₈, R₃₃₉, R₃₄₀, R₃₄₁, R₃₄₂, R₃₄₃, R₃₄₄, R₃₄₅, R₃₄₆, R₃₄₇, R₃₄₈, R₃₄₉, R₃₅₀, R₃₅₁, R₃₅₂, R₃₅₃, R₃₅₄, R₃₅₅, R₃₅₆, R₃₅₇, R₃₅₈, R₃₅₉, R₃₆₀, R₃₆₁, R₃₆₂, R₃₆₃, R₃₆₄, R₃₆₅, R₃₆₆, R₃₆₇, R₃₆₈, R₃₆₉, R₃₇₀, R₃₇₁, R₃₇₂, R₃₇₃, R₃₇₄, R₃₇₅, R₃₇₆, R₃₇₇, R₃₇₈, R₃₇₉, R₃₈₀, R₃₈₁, R₃₈₂, R₃₈₃, R₃₈₄, R₃₈₅, R₃₈₆, R₃₈₇, R₃₈₈, R₃₈₉, R₃₉₀, R₃₉₁, R₃₉₂, R₃₉₃, R₃₉₄, R₃₉₅, R₃₉₆, R₃₉₇, R₃₉₈, R₃₉₉, R₄₀₀, R₄₀₁, R₄₀₂, R₄₀₃, R₄₀₄, R₄₀₅, R₄₀₆, R₄₀₇, R₄₀₈, R₄₀₉, R₄₁₀, R₄₁₁, R₄₁₂, R₄₁₃, R₄₁₄, R₄₁₅, R₄₁₆, R₄₁₇, R₄₁₈, R₄₁₉, R₄₂₀, R

R_1 represents a hydrogen atom, a C_1 alkyl group (e.g. methyl or ethyl) or a phenyl or phenyl (C_6), 45 alkyl group in which the phenyl ring is unsubstituted or substituted by a C_1 alkoxy group (e.g. methoxy) or by the group $-CONH_2$;

R_1 and R_2 , each represents a hydrogen atom or a C_{1-6} alkyl group (e.g. methyl); and m is zero or 1;

and physiologically acceptable salts and solvates (e.g. hydrates) thereof.

50 In the compounds of formula (Ia) it is preferred that the total number of carbon atoms in R_{11} and R_{12} does not exceed two, and most preferably R_{11} and R_{12} each represent a methyl group. In compounds (Ia) m preferably represents zero.

Preferred compounds according to the invention include:

Preferred compounds according to the invention include:

55 (E)-2-[3-[2-(dimethylamino)ethyl]-1*H*-indol-5-yl]-*N*-(2-phenylethyl)-ethanesulphonamide;
(E)-2-[3-[2-(dimethylamino)ethyl]-1*H*-indol-5-yl]-*N*-[(4-methoxyphenyl)methyl]ethanesulphonamide;
(E)-2-[3-[2-(dimethylamino)ethyl]-1*H*-indol-5-yl]-*N*-[4-(methoxyphenyl)ethyl]ethanesulphonamide;

and the physiologically acceptable salts and solvates (e.g. hydrates) of these compounds.

Suitable physiologically acceptable salts of the indoles of general formula (I) include acid addition salts formed with inorganic or organic acids, for example hydrochlorides, hydrobromides, sulphates, nitrates, 60 phosphates, tartrates, citrates, fumarates, maleates, succinates, and sulphonates e.g. mesylates. Other

It will be appreciated that the invention extends to other physiologically acceptable equivalents of the compounds according to the invention, i.e. physiologically acceptable compounds which are converted *in vivo* into the parent compound. Examples of such equivalents include physiologically acceptable, metabolically labile N-acyl derivatives.

Compounds of the invention potently and selectively constrict the carotid arterial bed of the anaesthetised dog, whilst having a negligible effect on blood pressure. The selective vasoconstrictor action of compounds of the invention has been demonstrated *in vitro*.

Compounds of general formula (I) are useful in treating and/or preventing pain resulting from dilation of the cranial vasculature, in particular migraine and related disorders such as cluster headache.

Compounds of general formula (I') are preferred by virtue of their vasoconstrictor activity.

The invention also provides a pharmaceutical composition adapted for use in human medicine which comprises at least one compound according to the invention or a physiologically acceptable salt or solvate (e.g. hydrate) thereof and formulated for administration by any convenient route. Such compositions may be formulated in conventional manner using one or more pharmaceutically acceptable carriers or excipients.

Thus the compounds according to the invention may be formulated for oral, buccal, parenteral or rectal administration or in a form suitable for administration by inhalation or insufflation.

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropylmethylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium phosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium starch glycolate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g. sorbitol syrup, methyl cellulose or hydrogenated edible fats); emulsifying agents (e.g. lecithin or acacia); non-aqueous vehicles (e.g. almond oil, oily esters or ethyl alcohol); and preservatives (e.g. methyl or propyl-*p*-hydroxybenzoates or sorbic acid). The liquid preparations may also contain conventional buffers, flavouring, colouring and sweetening agents as appropriate.

For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

The compounds of the invention may be formulated for parenteral administration by injection or continuous infusion. Formulations for injection may be presented in unit dosage form e.g. in ampoules or in multi-dose containers, with an added preservative.

The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents, and/or agents to adjust the tonicity of the solution. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

The compounds of the invention may also be formulated in rectal compositions such as suppositories or retention enemas, e.g. containing conventional suppository bases such as cocoa butter or other glycerides.

For administration by inhalation the compounds according to the invention are conveniently delivered in the form of an aerosol spray presentation from pressurised packs, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas, or from a nebuliser. In the case of pressurised aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

A proposed dose of the compounds of the invention for oral, parenteral, buccal or rectal administration to man of average bodyweight e.g. about 70kg) for the treatment of migraine is 0.1 to 100mg of the active ingredient per unit dose which could be administered, for example, up to 8 times per day, more usually 1 to 4 times per day. It will be appreciated that it may be necessary to make routine variations to the dosage depending on the age and weight of the patient as well as the severity of the condition to be treated.

For oral administration a unit dose will preferably contain from 0.5 to 50mg e.g. 2 to 40mg of the active ingredient. A unit dose for parenteral administration will preferably contain 0.2 to 5mg of the active ingredient.

Aerosol formulations are preferably arranged so that each metered dose or 'puff' delivered from a pressurised aerosol contains 0.2 to 2mg of a compound of the invention and, each dose administered via capsules or cartridges in an inhaler or insufflator contains 0.2 to 20mg. The overall daily dose by inhalation will be within the range 1mg to 100mg. Administration may be several times daily, for example from 2 to 8 times, giving for example 1, 2 or 3 doses each time.

The compounds of the invention may, if desired, be administered in combination with one or more other therapeutic agents, such as analgesics, anti-inflammatory agents and anti-nauseants.

In addition to their vasoconstrictor activity, compounds of general formula (I) are also useful as intermediates for the preparation of further indole derivatives. Thus, compounds of formula (I) may be reduced to give compounds of formula (II):



(II)

wherein R_1 , R_2 , R_3 , R_4 , and Alk are as previously defined, and A represents an alkyl chain containing two to five carbon atoms.

Compounds of formula (II) wherein R_1 represents a substituted phenyl or substituted phenyl (C_{6H_5}) alkyl group are described in our published European Application No. 147107. Compounds of formula (II) wherein R_1 represents a hydrogen atom, a C_{1-3} alkyl, C_{2-3} alkenyl, or C_6 cycloalkyl group, or an unsubstituted phenyl or phenyl (C_{6H_5}) alkyl group are disclosed in our published UK Application No. 2150932A.

The reduction of compounds of formula (I) to give compounds of formula (II) may be effected by methods well known in the art.

Thus, for example, a compound of formula (I) may be reduced by catalytic hydrogenation, using a heterogeneous or homogeneous catalyst. Heterogeneous catalysts which may be employed include Raney nickel; nickel reduced with sodium borohydride; and noble metal catalysts such as platinum, platinum oxide, palladium, palladium oxide, rhodium or ruthenium, which may be supported for example on charcoal, kieselguhr or alumina. In the case of Raney nickel, hydrazine may also be used as the source of hydrogen. Examples of homogeneous catalysts include chlorotris (triphenylphosphine)rhodium and pentacyano cobaltate. The catalytic hydrogenation may conveniently be carried out in a solvent such as an alcohol e.g. ethanol; an ether, e.g. dioxan or tetrahydrofuran, an amide, e.g. dimethylformamide; or an ester e.g. ethyl acetate, and at a temperature of from -10 to $+50^\circ\text{C}$, preferably -5 to $+30^\circ\text{C}$. The reaction may conveniently be effected at atmospheric pressure, but higher pressures, e.g. up to 5 atmospheres, may be employed.


The compounds of the present invention may also be reduced with other reducing agents such as sodium in ethanol, or sodium and *t*-butylalcohol in hexamethylphosphoramide, at a temperature of from 0 to 120°C .

The following compounds of general formula (II) which may be prepared from the corresponding compounds of formula (I) according to the above-described process, are novel compounds and constitute a further feature of the present invention:

- 3-[2-(dimethylamino)ethyl]-N-methyl-1*H*-indole-5-propanesulphonamide;
- 3-[2-(dimethylamino)ethyl]-N,N-dimethyl-1*H*-indole-5-ethanesulphonamide;
- 3-[2-(dimethylamino)ethyl]-N-(2-phenylethyl)-1*H*-indole-5-ethanesulphonamide;
- 3-[2-(dimethylamino)ethyl]-N-(1-methylethyl)-1*H*-indole-5-ethanesulphonamide;
- 3-[2-(dimethylamino)ethyl]-N-ethyl-1-indole-5-ethanesulphonamide;
- 3-[2-(dimethylamino)ethyl]-N-phenyl-1*H*-indole-5-ethanesulphonamide;
- and, ethyl
- N-cyclopentyl-3-[2-(dimethylamino)-1*H*-indole-5-ethane sulphonamide.


According to another aspect of the invention, compounds of general formula (I) and their salts and solvates may be prepared by the general methods outlined hereinafter. In the following processes R_1 , R_2 , R_3 , R_4 , A' , Alk, m and n are as defined for the general formula (I) unless otherwise specified.

According to a general process (A), compounds of general formula (I) may be prepared by reacting an indole of general formula (III):



(III)

wherein X represents a leaving atom or group such as a halogen atom, e.g. a bromine or iodine atom with an alkene of formula (IV):



(IV)

wherein $A'-CH_2$ represents a C_{1-5} alkenyl chain.

The reaction will generally be effected in the presence of a palladium catalyst and a base. The catalyst may be for example palladium on charcoal or a palladium salt. Palladium salts which may be employed as catalysts include salts of organic acids, e.g. acetates, and salts of inorganic acids e.g. chlorides or bromides. The base may be for example a tertiary nitrogen base such as triethylamine, or tri-*n*-butylamine or an alkali metal carbonate, e.g. sodium carbonate. The reaction may optionally be carried out in the presence of a phosphine, for example a triarylphosphine such as triphenylphosphine or tri-*o*-tolylphosphine. A phosphine should be present when the process is effected with a compound of formula (III) wherein X represents a bromine atom. The reaction is conveniently carried out using a small excess of

the alkene (V) with respect to the indole (III). It is generally preferred that an excess of the base (e.g. ca. 3 equivalents) and, when present an excess of the phosphine (e.g. ca. 2 equivalents) are also employed.

General process (A) may be effected in the presence or absence of solvent. An anhydrous or aqueous reaction medium comprising one or more solvents may be employed. Suitable solvents include nitriles, e.g. acetonitrile; alcohols e.g. methanol or ethanol; amides e.g. dimethylformamide, N-methylpyrrolidone or hexamethylphosphoramide; and water. The reaction may conveniently be carried out at a temperature of from 25 to 200°C, preferably 75 to 150°C.

In the compounds of formula (IV) the moiety $-A^1-CH_2-$ preferably represents the group $-(CH_2)_m-CH_2-$, wherein m is zero or an integer from 1 to 3.

10 It will be appreciated that the compounds of formula (I) prepared by general process (A) will be those in which n is zero.

According to another general process (B) compounds of general formula (II) may be prepared by reacting an aldehyde of formula (V):




(wherein A' represents a bond or a C₁₋₄ alkyl chain) with a reagent serving to form the group R₁R₂NSO₂A'-.

A suitable reagent serving to form the group R,R',NSO_2A'' may be, for example, a phosphorus ylide of general formula (VI):



(wherein A' represents a bond or a C₁, alkyl chain such that the total number of carbon atoms in A' and A' does not exceed 3, and R₁ is an alkyl, e.g. methyl, or aryl, e.g. phenyl or tolyl group) or a phosphonate ester of general formula (VII):




(wherein A¹ represents an alkyl chain containing 1 to 4 carbon atoms, such that the total number of carbon atoms in A¹ and A² does not exceed 4, and R₁ represents an alkyl e.g. methyl; aryl; aryl e.g. phenyl or aralkyl e.g. benzyl group).

The reaction with an ylide of formula (VI) may conveniently be effected in an anhydrous reaction medium which may comprise one or more organic solvents. Solvents which may be employed include amides e.g. dimethylformamide; ethers, e.g. acyclic ethers such as diethyl ether and cyclic ethers such as tetrahydrofuran; and hydrocarbons e.g. xylene or toluene. The reaction may conveniently be conducted at a temperature of from -70 to $+150^{\circ}\text{C}$.

45 at a temperature of from -70 to $+150^{\circ}\text{C}$.
A phosphonate ester of formula (VII) will preferably be reacted with an aldehyde of general formula (V) in the presence of a base, for example a metal hydride, such as sodium or potassium hydride; a metal amide such as sodium amide; an alkali metal alkoxide, such as potassium *t*-butoxide; or an organolithium base, such as butyllithium. The reaction may be conveniently effected in an organic reaction medium, which may comprise one or more solvents, and at a temperature in the range -70 to $+150^{\circ}\text{C}$. Suitable solvents include amides, ethers and hydrocarbons, such as those mentioned above for the reaction with an ylide of formula (VI).

Phosphorus ylides of formula (VI) may be prepared by reaction of the corresponding phosphonium salt of formula (VIII):



(wherein A⁺ and R_n are as previously defined and E⁻ represents an anion, such as a halide ion, e.g. a chloride, bromide or iodide ion; or a sulphonate anion, e.g. methanesulphonate or p-toluene sulphonate) with a base. Bases which may be employed include organolithium compounds e.g. n-butyllithium and phenyllithium; metal hydrides, e.g. sodium hydride; metal amides, e.g. sodium amide; alkali metal alkoxides e.g. sodium or potassium methoxide, ethoxide or t-butoxide; and alkali metal carbonates e.g. sodium carbonate. The formation of the phosphorus ylide may be effected in an organic solvent or mixture of solvents, for example as described for general process (B).

In a particular embodiment of general process (B), an aldehyde of general formula (V) may be reacted directly with a phosphonium salt of formula (VIII) in the presence of a base, using the reaction conditions described above for the reaction of an aldehyde (V) with an ylide of general formula (VI).


Compounds of formula (V) may be prepared by reacting a corresponding nitrile of formula (IX):



(wherein A' is as previously defined for general formula (V)) with a reducing agent such as di-isobutylaluminum hydride, in a solvent such as tetrahydrofuran, followed by hydrolysis, which may be effected for example by the addition of water. The reaction may be effected at a temperature of -70 to 30°C .

Compounds of formula (IX) may be prepared by cyclisation of a corresponding hydrazone, in an analogous manner to process (D) described hereinafter.


Compounds of general formula (I) may also be prepared according to a further general process (C), which comprises elimination of HX' from a compound of formula (X):



(wherein A' represents a C_1 alkyl chain substituted by a leaving atom or group, X' , for example a halogen atom, a hydroxy group or an acyloxy group).

The group A' may for example be represented by the formula

X'



When X' in the group A' represents a halogen atom, this may be, for example, bromine or chlorine. An acyloxy group X' may be derived from a carboxylic or sulphonic acid, such as an acetoxy, chloroacetoxy, p-nitrobenzoyloxy, p-toluenesulphonyloxy or methanesulphonyloxy group.


When X' represents a halogen atom or an acyloxy group, the elimination may be effected thermally, e.g. at a temperature of 30 to 200°C , or using a base such as an alkali metal alkoxide, e.g. sodium or potassium ethoxide or t-butoxide; an alkali metal hydroxide, e.g. sodium or potassium hydroxide; or a tertiary amine base e.g. triethylamine. The reaction with a base may be effected in an organic reaction medium, at a temperature in the range -10 to $+150^\circ\text{C}$. Solvents which may be employed include alcohols e.g. ethanol or t-butanol; amides e.g. dimethylformamide; sulphoxides e.g. dimethylsulphoxide; halogenated hydrocarbons e.g. methylene chloride; ketones e.g. acetone and esters e.g. ethyl acetate, as well as mixtures of such solvents.

When X' represents a hydroxy group compounds of formula (X) may be heated with an acid such as sulphuric or phosphoric acid, to give a compound of formula (I).

Compounds of the formula (X) wherein X' represents an acyloxy group may be prepared for example by reacting the corresponding compound wherein X' is a hydroxyl group, with an appropriate acylating agent, such as an acid halide e.g. methanesulphonyl chloride. Compounds of formula (X) wherein X' represents a hydroxyl group may also be used to prepare corresponding compounds wherein X' is a halogen atom, for example, by reaction with the appropriate phosphorus trihalide.

Compounds of formula (X) wherein X' represents a hydroxyl group may themselves be prepared by condensing an aldehyde of general formula (V) with an appropriate alkane sulphonamide in the presence of a base such as n-butyllithium or lithium di-isopropylamide at temperatures of from -80 to -10°C .

A further general process (D) for preparing compounds of general formula (XI):



wherein Q is the group NR_3R_4 (or a protected derivative thereof) or a leaving atom or group such as a halogen atom (e.g. chlorine or bromine) or an acyloxy group, for example a carboxylic or sulphonic acyloxy group such as acetoxy, chloroacetoxy, dichloroacetoxy, trifluoroacetoxy, p-nitrobenzoyloxy, p-tolu-

enesulphonloxy or methanesulphonyloxy group.

The reaction may conveniently be effected in aqueous or non-aqueous reaction media, and at temperatures of from 20 to 200°C, preferably 50 to 125°C.

Particularly convenient embodiments of the process are described below.

- 5 When Q is the group NR₂R₄ (or a protected derivative thereof) the process is desirably carried out in the presence of polyphosphate ester in a reaction medium which may comprise one or more organic solvents, preferably halogenated hydrocarbons such as chloroform, dichloromethane, dichloroethane, dichlorodifluoromethane, or mixtures thereof. Polyphosphate ester is a mixture of esters which may be prepared from phosphorus pentoxide, diethylether and chloroform according to the method described in 10 'Reagents for Organic Synthesis', (Fieser and Fieser, John Wiley and Sons 1967).
- Alternatively the cyclisation may be carried out in an aqueous or non-aqueous reaction medium, in the presence of an acid catalyst. When an aqueous medium is employed this may be an aqueous organic solvent such as an aqueous alcohol (e.g. methanol, ethanol or isopropanol) or an aqueous ether (e.g. dioxan or tetrahydrofuran) as well as mixtures of such solvents and the acid catalyst may be, for exam- 15 ple, an inorganic acid such as concentrated hydrochloric or sulphuric acid or an organic acid such as acetic acid. (In some cases the acid catalyst may also act as the reaction solvent). In an anhydrous reaction medium, which may comprise one or more ethers (e.g. as previously described) or esters (e.g. ethyl acetate), the acid catalyst will generally be a Lewis acid such as boron trifluoride, zinc chloride or magnesium chloride.
- 20 When Q is a leaving atom or group such as a chlorine or bromine atom the reaction may be effected in an aqueous organic solvent, such as an aqueous alcohol (e.g. methanol, ethanol or isopropanol) or an aqueous ether (e.g. dioxan or tetrahydrofuran) in the absence of an acid catalyst, conveniently at a temperature of from 20 to 200°C, preferably 50 to 125°C. This process results in the formation of a compound of formula (I) wherein R₁ and R₂ are both hydrogen atoms.
- 25 According to a particular embodiment of this process compounds of formula (I) may be prepared directly by the reaction of a compound of general formula (XII):



or a salt thereof, with a compound of formula (XIII):



(XIII)

- 40 wherein Q is as defined above or a salt or protected derivative thereof (such as an acetal or ketal e.g. formed with an appropriate alkyl orthoformate or diol, or protected as a bisulphite addition complex) using the appropriate conditions as described above for the cyclisation of compounds of general formula (XI). It will be appreciated that in this embodiment of the cyclisation process (D) a compound of general formula (XI) is formed as an inter- 45 mediate, and may be reacted *in situ* to form the desired compound of general formula (I).
- Compounds of general formula (XI) may, if desired, be isolated as intermediates during the process for the preparation of compounds of formula (I) wherein a compound of formula (XII), or a salt or protected derivative thereof, is reacted with a compound of formula (XIII) or a salt or protected derivative thereof, in a suitable solvent, such as an aqueous alcohol (e.g. methanol) at a temperature of, for example, 20 to 50 30°C. If an acetal or ketal of a compound of formula (XIII) is used, it may be necessary to carry out the reaction in the presence of an acid (for example, acetic or hydrochloric acid).
- Compounds of general formula (XII) may be prepared for example from the corresponding nitro compounds, using conventional procedures.

- A further general process (E) for preparing compounds of general formula (I) involves reacting a compound of general formula (XIV):



(wherein Y is a readily displaceable atom or group) or a protected derivative thereof, with an amine of formula R₃R₄NH.

- 65 The displacement reaction may conveniently be carried out on those compounds of formula (XIV)


wherein the substituent Y is a leaving atom or group such as a halogen atom (e.g. chlorine, bromine or iodine) or a group OR, where OR, is, for example, an acyloxy group which may be derived from a carboxylic or sulphonic acid, such as an acetoxy, chloroacetoxy, dichloroacetoxy, trifluoroacetoxy, p-nitrobenzyloxy, p-toluenesulphonyloxy or methanesulphonyloxy group.

5 The displacement reaction may be conveniently effected in an inert organic solvent (optionally in the presence of water), examples of which include alcohols, e.g. ethanol; cyclic ethers, e.g. dioxan or tetrahydrofuran; acyclic ethers e.g. diethylether; esters, e.g. ethyl acetate; amides, e.g. N,N-dimethylformamide; and ketones e.g. acetone or methylethylketone, at a temperature of from -10 to $+150^{\circ}\text{C}$, preferably 20 to 50°C .

10 The compounds of general formula (XIV) wherein Y is a halogen atom may be prepared by conventional procedures in which a hydrazine of general formula (XII) is reacted with an aldehyde or ketone (or a protected derivative thereof) of formula (XIII) in which Q is a halogen atom, in an aqueous alcohol (e.g. methanol) containing an acid (e.g. acetic or hydrochloric acid). Compounds of formula (XIV) wherein Y is the group OR, may be prepared from the corresponding compound wherein Y is a hydroxyl group by


15 acylation with the appropriate activated species (e.g. anhydride or sulphonyl chloride) using conventional techniques. The intermediate alcohol may be prepared by cyclisation of a compound of formula (XI) wherein Q is a hydroxyl group (or a protected derivative thereof) under standard conditions.

Compounds of general formula (I) may also be prepared by another general process (F) which comprises reacting an indole of general formula (XV):



(XV)

wherein Z represents a leaving atom or group with a compound of general formula (XVI):



(XVI)

Examples of suitable leaving atoms or groups Z in the compound of general formula (XVI) include a halogen atom (e.g. a fluorine, chlorine or bromine atom) or a group OR, where R, represents a hydrocar-
 35 byl group such as an aryl group, e.g. phenyl. The aryl group may be unsubstituted or substituted by one or more substituents such as halogen atoms; or nitro; cyano; amino; alkyl e.g. methyl; alkoxy e.g. methoxy; acyl e.g. acetyl and alkoxy carbonyl e.g. ethoxycarbonyl groups. The leaving group represented by Z is preferably a phenoxy group.


The reaction is conveniently carried out in the presence of a solvent and may be effected in an aqueous or non-aqueous reaction medium.

The reaction medium may thus comprise one or more organic solvents, such as ethers, e.g. dioxan or tetrahydrofuran; amides e.g. N,N-dimethylformamide or N-methylpyrrolidone; alcohols e.g. methanol or ethanol; esters e.g. ethyl acetate, nitriles e.g. acetonitrile; halogenated hydrocarbons e.g. dichloromethane; and tertiary amines e.g. triethylamine or pyridine, optionally in the presence of water. In some
 45 cases the amino of formula (XVI) may itself serve as the solvent.

If desired the aminolysis may be effected in the presence of a base, such as a tertiary amine (e.g. triethylamine or pyridine); an alkoxide (e.g. potassium t-butoxide); a hydride (e.g. sodium hydride); or an alkali metal carbonate (e.g. sodium carbonate).

The reaction may conveniently be effected at a temperature of from -20°C to $+150^{\circ}\text{C}$.

50 The starting materials of general formula (XV) may be prepared for example by cyclisation of a compound of general formula (XVII):



(XVII)

(wherein Z and Q are as previously defined).

The cyclisation may be effected in an analogous manner to the general process (D), described above.

60 According to a further general process (G) a compound of formula (I) according to the invention, or a salt or protected derivative thereof, may be converted into another compound of formula (I) using conventional procedures.

For example, a compound of general formula (I) wherein one or more of R₁, R₂, R₃, and R₄ are alkyl groups may be prepared from the corresponding compounds of formula (I) wherein one or more of R₁, R₂, R₃, and R₄ represent hydrogen atoms, by reaction with a suitable alkylating agent such as a compound
 65

of formula R₁L (where R₁ represents the desired R₁, R₂, R₃ or R₄ group and L represents a leaving atom or group such as a halogen atom or a tosylate group) or a sulphate (R₁)₂SO₄. Thus, the alkylating agent may be for example an alkyl halide (e.g. methyl or ethyl iodide), alkyl tosylate (e.g. methyl tosylate) or dialkylsulphate (e.g. dimethylsulphate).

- 5 The alkylation may conveniently be carried out in an inert organic solvent such as an amide (e.g. dimethylformamide), an ether (e.g. tetrahydrofuran) or an aromatic hydrocarbon (e.g. toluene) preferably in the presence of a base. Suitable bases include, for example, alkali metal hydrides such as sodium or potassium hydride; alkali metal amides such as sodium amide; alkali metal carbonates such as sodium carbonate; alkali metal alkoxides such as sodium or potassium methoxide, ethoxide or t-butoxide; and
10 tetrabutylammonium fluoride. When an alkyl halide is employed as the alkylating agent the reaction may also be carried out in the presence of an acid scavenging agent such as propylene or ethylene oxide. The reaction may be conveniently effected at a temperature of from -20° to 100°C.

- Compounds of formula (I) wherein R₁ represents an alkenyl group, R₂ represents an alkenyl, phenylalkyl or cycloalkyl group and/or one or both of R₃ and R₄ represents propenyl may be prepared similarly, using
15 an appropriate compound of formula R₁L or (R₁)₂SO₄.

According to another general process (H), a compound of general formula (I) according to the invention, or a salt thereof may be prepared by subjecting a protected derivative of general formula (I) or a salt thereof to reaction to remove the protecting group or groups.

- Thus, at an earlier stage in the reaction sequence for the preparation of a compound of general formula (I) or a salt thereof it may have been necessary or desirable to protect one or more sensitive groups
20 in the molecular to avoid undesirable side reactions. For example it may be necessary to protect the group NR₁R₂, wherein R₃ and/or R₄ represents hydrogen, by protonation or with a group easily removable at the end of the reaction sequence. Such groups may include, for example, aralkyl groups, such as diphenylmethyl or triphenylmethyl; or acyl groups such as N-benzoyloxycarbonyl or t-butoxycarbonyl or
25 phthaloyl.

Subsequent cleavage of the protecting group or groups may be achieved by conventional procedures. Thus an aralkyl group such as triphenylmethyl may be cleaved by treatment with dilute acid e.g. dilute hydrochloric acid; and an acyl group such as N-benzoyloxycarbonyl may be removed by hydrolysis with, for example, hydrogen bromide in acetic acid.

- 30 The phthaloyl group may be removed by hydrazinolysis (e.g. by treatment with hydrazine hydrate) or by treatment with a primary amine (e.g. methylamine).

- As will be appreciated, in some of the general processes (A) to (G) described previously it may be necessary or desirable to protect any sensitive groups in the molecular as just described. Thus, a reaction step involving deprotection of a protected derivative of general formula (I) or a salt thereof may be carried out subsequent to any of the previously described processes (A) to (G).
35

Thus, according to a further aspect of the invention, the following reactions in any appropriate sequence may if necessary and/or desired be carried out subsequent to any of the processes (A) to (G):

- (i) removal of any protecting groups; and
(ii) conversion of a compound of general formula (I) or a salt thereof into a physiologically acceptable
40 salt or solvate (e.g. hydrate) thereof.

Where it is desired to isolate a compound of the invention as a salt, for example as an acid addition salt, this may be achieved by treating the free base of general formula (I), with an appropriate acid, preferably with an equivalent amount or with creatinine sulphate in a suitable solvent (e.g. aqueous ethanol).

- The starting materials or intermediate compounds for the preparation of the compounds according to this invention may be prepared for example by analogous methods to those described in UK Published
45 Patent Application No. 2035310 and 2124210.

As well as being employed as the last main step in the preparative sequence, the general methods indicated above for the preparation of the compounds of the invention may also be used for the introduction of the desired groups at an intermediate stage in the preparation of the required compound.

- 50 Thus, for example, the required group at the 5-position may be introduced before or after cyclisation to form the indole nucleus. It should therefore be appreciated that in such multi-stage processes, the sequence of reactions should be chosen in order that the reaction conditions do not affect groups present in the molecule which are desired in the final product.

The invention is further illustrated by the following Examples. All temperatures are in °C. chromatography was carried out either in the conventional manner using silica gel (Merck, Kieselgel 60, Art. 7734) or
55 by flash Chromatography (W.C. Still, M. Kahn and A. Mitra, J. Org. Chem. 2933, 43, 1978) on silica (Merck 9385) and thin layer chromatography (t.l.c.) on silica (Machery-Nagel, Polygram) except where otherwise stated. The following abbreviations define the eluent used for chromatography and t.l.c.

- (A) Methylene chloride-ethanol-0.88 ammonia 50:8:1
60 (B) Methylene chloride-ethanol-0.88 ammonia 100:8:1
(C) Methylene chloride-ether 1:1
(D) Methylene chloride-ethanol-0.88 ammonia 200:8:1
(E) Cyclohexane-ether 2:1
(F) Cyclohexane-ether 1:1
65 Intermediates were routinely checked for purity by t.l.c. employing u.v. light for detection and spray

reagents such as potassium permanganate (KMnO_4). In addition indolic intermediates were detected by spraying with aqueous ceric sulphate (Ce^{IV}) and tryptamines by spraying with a solution of iodoplatinic acid (IPA) or ceric sulphate.

Proton (^1H) nuclear magnetic resonance (n.m.r.) spectra were obtained either at 90MHz using a Varian EM 390 instrument or at 250MHz using a Bruker AM or WM 250 instrument. s = singlet, d = doublet t = triplet, m = multiplet and q = quartet.

Reactivials are 4ml stout-walled glass vials with a screw cap and teflon-faced disc, supplied by Pierce and Warriner (UK) Ltd.

10 Preparation 1

N-Methyl-2-propenesulphonamide

Dry methylamine gas was bubbled through a solution of 2-propenesulphonyl chloride (5.24g) in dry ether (50ml) whilst maintaining the internal temperature at -78° . After 30 min, the flow of methylamine was stopped and the reaction mixture stirred at -78° for an additional period of 45 min. On allowing to warm to ambient temperature, water (100ml) was added and the reaction mixture acidified (5N HCl; to pH1). The ethereal layer was separated and the aqueous phase extracted with dichloromethane (5 x 100ml). The combined organic extracts were dried (Na_2SO_4) and concentrated *in vacuo* to afford the *title compound* as an oil (1.41g)

T.l.c. (C) Rf 0.65 N.m.r. $\delta(\text{CDCl}_3)$ 2.80(3H,d, SO_2NHMe), 3.72 (2H,d, $\text{CH}_2\text{SO}_2\text{NH}$), 5.3-6.2 (3H,m, $\text{CH}_2=\text{CH}$)

20 Preparation 2

N-(2-Phenylethyl)ethanesulphonamide

2-Chloroethanesulphonyl chloride (8.16g) was dissolved in benzene (30ml), the solution cooled to 5° , stirred well and treated with 2-phenylethylamine (20g) in benzene (12.5ml). The mixture was stirred for a further 1h, then washed with dilute hydrochloric acid (25ml) and sodium hydrogen carbonate (8%, 60ml) and dried to give an oil (10.3g). This oil was distilled to give the product as an oil (2.2g) which was then further purified by flash chromatography (E) to give the *title compound* (1.63g) as an oil. T.l.c. (F) Rf 0.3 (KMnO_4).

30 Preparation 3

N-Cyclopentylathanesulphonamide

A mixture of cyclopentylamine (8.5g) and triethylamine (27.8ml) in ether (50ml) was added dropwise over 0.5h to a stirred solution of 2-chloroethanesulphonyl chloride (16.2g) in anhydrous ether (200ml) at ca. 65° . The mixture was allowed to reach 15° over a period of 1h, the suspension filtered and the filtrate concentrated *in vacuo* to give an oil (10.5g), which was purified by chromatography (dichloromethane). A portion of the resulting oil (1.5g) was distilled at 135/6 mmHg to give the *title compound* (1.2g) as an oil.

T.l.c. (dichloromethane) Rf 0.5 (KMnO_4)

40 Preparation 4

N-[4-Methoxyphenyl)methyl]athanesulphonamide

A cold solution of 4-methoxybenzylamine (2g) and triethylamine (2.8ml) in dry dichloromethane (20ml) at -78° was transferred under nitrogen to a solution of 2-chloroethanesulphonyl chloride (4.9g) in dry dichloromethane (20ml) at -78° . The mixture was stirred for 4h whilst warming to room temperature and then refrigerated overnight. Water (ca 100ml) was added and the organic layer separated. This was washed with hydrochloric acid (2N, 50ml), water (50ml) and brine (50ml), dried (MgSO_4) and evaporated under reduced pressure. The residue was purified by chromatography (dichloromethane) to give the *title compound* as a powder (2g) m.p. $68-69^\circ$.

50 Preparation 5

4-[[[Ethenylsulphonyl]amino]methyl]benzamide

A solution of 4-aminomethylbenzamide (0.58g) and triethylamine (1.1ml) in dimethylformamide (DMF; 6ml) was added to a solution of 2-chloroethanesulphonyl chloride (0.63g) in DMF (4ml) at -60° under nitrogen over 30 mins. The mixture was allowed to warm to room temperature and stirred for 18h. The mixture was evaporated to give a semi-solid (2.78g) which was purified by column chromatography (D) to give the *title compound* as a solid (0.54g) m.p. $142-4^\circ$.

Assay Found : C,50.0; H,5.3; N,11.5. $\text{C}_{10}\text{H}_{11}\text{N}_2\text{O}_3\text{S}$ requires C,50.0; H,5.0; N,11.7%.

Preparation 6

60 5-Iodo-N,N-dimethyl-1H-indole-3-ethanamine oxalate

(1) 4-(dimethylamino)butanone (4-Iodophenyl)hydrazine

A solution of 4-Iodophenylhydrazine (2g) in water (70ml) and 2N hydrochloric acid (4ml) was stirred at room temperature with 4-dimethylaminobutanal, diethyl acetal (2.6g) for 3h. The resulting solution was partitioned between sodium bicarbonate (50ml) and ethyl acetate (2x50ml). The combined organic extracts were dried (Na_2SO_4) and evaporated *in vacuo* to give an oil (2.3g), which was used directly in the

next stage.

T.l.c. (B) Rf 0.3.

5 (ii) 5-Iodo-N,N-dimethyl-1H-indole-3-ethanamine oxalate 5

A solution of the product of Stage (i) (2.3g) and polyphosphate ester (40g) in chloroform (80ml) was refluxed for 5 min. The solution was added to ice (300g), stirred for 20min, poured into 2N aqueous sodium carbonate (100ml) and extracted with chloroform (2x100ml). The combined organic extracts were dried (Na₂SO₄) and evaporated *in vacuo*. The resulting oil was purified by flash chromatography (B) to

10 give pure free base as a solid. A solution of the base (0.92g) in ethanol (20ml) was added to oxalic acid (0.28g) in methanol (5ml) and the *title compound* precipitated. m.p. 176-177°.

T.l.c. (B) Rf 0.3.

Analysis Found: C,41.6; H,4.2; N,6.9. C₁₁H₁₁N₂O₂ requires C,41.3; H,4.1; N,6.55%.

15 Example 1 15

(E)-2-[3-[2-(Dimethylamino)ethyl]-1H-indol-5-yl]-N-methyl-2-propenesulphonamide oxalate

A mixture of the product of Preparation 1 (247mg), 5-bromo-N,N-dimethyl-1H-indole-3-ethanamine oxalate (650mg), palladium acetate (8.3mg), tri-ortho-tolylphosphine (26.3mg) and triethylamine (1.05ml) in acetonitrile (3ml) was heated in a "reactivial" at 105-110° for a period of 24h. On cooling to ambient

20 temperature, the reaction mixture was poured into water (20ml) and the emulsion extracted with ethyl acetate (3 x 50ml). The combined organic extracts were dried (Na₂SO₄) and concentrated *in vacuo*. Flash chromatography (B) of the residue afforded the free base as a foam (283mg). A filtered solution of the free base (272.5mg) in absolute ethanol (0.5ml) was added to a solution of oxalic acid (76.3mg) in absolute ethanol (0.75ml) from which a solid was deposited on scratching. The salt was filtered off (240mg),

25 washed with ether (20ml) dried and recrystallised from ethanol (20ml) to afford the *title compound* as a powder (98mg) m.p. 93-95°.

Analysis Found: C,52.4; H,6.5; N,10.2. C₁₁H₁₁N₂O₂S.C₂H₂O₄ requires C,52.5; H,6.1; N,10.2%.

N.m.r. δ (CD₃SOCD₃) includes 2.66(3H,s,SO,NHMe), 2.81(6H,s,NMe₂), 3.05-3.3(4H,m,CH, CH,N), 3.96(2H,d,SO,CH₂CH=CH), 6.15 (1H,dt,CH₂CH=CH), 6.88(1H,d,CH₂CH=CH), 7.2-7.7(4H,m,aromatic)

30

Example 2

The following compounds were prepared using a similar method to that in Example 1, the appropriate alkenesulphonamide and the reaction conditions shown in Table I.

(a)(E)-2-[3-[2-(Dimethylamino)ethyl]-1H-indol-5-yl]ethanesulphonamide oxalate

35 m.p. 192°(dec). Analysis Found: C,49.4; H,5.5; N,10.5. C₁₁H₁₁N₂O₂S.C₂H₂O₄.0.44H₂O requires C,49.1; H,5.6; N,10.7%. N.m.r. δ (CD₃SOCD₃) 2.85(6H,s,NMe₂), 3.1-3.35 (4H,m,CH₂CH₂N), 7.15(1H,d,SO,CH=CH), 7.43(1H,d,SO,CH=CH), 7.3-8.0(4H,m,aromatic)

(b)(E)-2-[3-[2-(Dimethylamino)ethyl]-1H-indol-5-yl]-N-methylethanesulphonamide oxalate

40 m.p. 189-190°. Analysis Found: C,50.95; H,6.2; N,10.45. C₁₁H₁₁N₂O₂S.C₂H₂O₄.0.21H₂O requires C,50.9; H,5.9; N,10.5%. N.m.r. δ (CD₃SOCD₃) 2.83(6H,s,NMe₂), 3.05-3.35 (4H,m,CH₂CH₂N), 7.01(1H,d,SO,CH=CH), 7.45(1H,d,SO,CH=CH), 7.3-8.0(5H,m,aromatic + NHSO₂)

(c)(E)-2-[3-[2-(Dimethylamino)ethyl]-1H-indol-5-yl]-N,N-dimethylethanesulphonamide oxalate

45 m.p. 136-138°. Analysis Found: C,51.6; H,6.0; N,9.5. C₁₁H₁₁N₂O₂S.C₂H₂O₄.0.033C₂H₂O requires C,51.4; H,6.4; N,9.6%. N.m.r. δ (CD₃SOCD₃) 2.75(6H,s,SO,NMe₂), 2.84(6H,s,NMe₂), 3.05-3.35(4H,m,CH₂CH₂N), 7.15(1H,d,SO,CH=CH), 7.51(1H,d,SO,CH=CH), 7.3-8.05(4H,m, aromatic).

(d)(E)-2-[3-[2-(Dimethylamino)ethyl]-1H-indol-5-yl]-N-(2-phenylethyl)ethanesulphonamide hemifumarate

50 m.p. 186-189°. Analysis Found: C,62.8; H,6.4; N,9.0. C₁₁H₁₁N₂O₂S.O.5C₄H₂O₄.0.013H₂O requires C,62.9; H,6.4; N,9.2%. N.m.r. δ (CD₃SOCD₃) 2.27(6H,s,NMe₂), 2.56(2H,m,CH₂), 2.75-2.9(4H,m,CH₂CH₂N and PhCH₂CH₂), 3.16(2H,m,CH₂NH), 6.9(1H,d,SO,CH=CH), 7.15-7.9(10H,m,aromatic + NHSO₂CH=CH).

(e)(E)-2-[3-[2-(Dimethylamino)ethyl]-1H-indol-5-yl]-N-(1-methylethyl)ethanesulphonamide oxalate

55 m.p. 125-129° Analysis Found: C,53.1; H,6.5; N,9.8. C₁₁H₁₁N₂O₂.C₂H₂O₄.0.12H₂O requires C,53.4; H,6.4; N,9.8%. N.m.r. δ (CD₃SOCD₃) 1.12(6H,d,CHMe₂), 2.79 (6H,s,NMe₂), 3.05-3.2(4H,m,CH₂CH₂N), 3.37(1H,m,CHMe₂), 7.02(1H,d,SO,CH=CH), 7.3-8.0(5H,m,aromatic + SO₂CH=CH).

(f)(E)-2-[3-[2-(Dimethylamino)ethyl]-1H-indol-5-yl]-N-ethylethanesulphonamide hemifumarate

60 m.p. 200-201°. Analysis Found: C,58.3; H,6.7; N,10.7. C₁₁H₁₁N₂O₂S.O.5C₄H₂O₄.0.15H₂O requires C,58.6; H,6.6; N,11.0%. N.m.r. δ (CD₃SOCD₃) 1.10(3H,t,SO,NHCH₂CH₃), 2.40 (6H,s,NMe₂), 2.7-3.0(6H,m,CH₂CH₂N and SO₂NHCH₂CH₃), 7.01(1H,d,SO,CH=CH), 7.25-7.95 (5H,m,aromatic + SO₂CH=CH)

(g)(E)-N-Cyclopentyl-2-[3-[2-(dimethylamino)ethyl]-1H-indol-5-yl]ethanesulphonamide oxalate

65 m.p. 202-203° Analysis Found: C,55.1; H,6.5; N,9.1. C₁₁H₁₁N₂O₂S.C₅H₉O₄.0.27H₂O requires C,55.3; H,6.4;

N,9,2%. N.m.r. δ (CD₃SOCD₃) 1.4-1.9(8H,m,cyclopentylmethylene protons), 2.83(6H,s,NMe₂), 3.05-3.35(4H,m,CH₂CH₂N), 3.55(1H,m,SO₂NHCH₂), 7.02(1H,d,SO₂CH=CH), 7.3-8.0(5H,m,aromatics + SO₂CH=CH)

(h)(E)-2-[3-[2-(Dimethylamino)ethyl]-1H-indol-5-yl]-N-phenylthene sulphonamide hemioxalate.
5 m.p. 203-205° (d) Analysis Found: C,59.4; H,5.7; N,9.4; C₂₀H₂₃N₃O₄S,0.5C₂H₂O₄,0.025 EtOH requires C,59.1; H,6.1; N,9.6%. N.m.r. δ (CD₃SOCD₃) 2.53(6H,s,NMe₂), 2.8-3.0(4H,m,CH₂CH₂N), 7.0-8.0(12H,m,aromatics + SO₂CH=CH + 2NH

(i)(E)-2-[3-[2-(Dimethylamino)ethyl]-1H-indol-5-yl]-N-[(4-methoxyphenyl)methyl]ethenesulphonamide oxalate
10 late m.p. 166-169°. Analysis Found: C,57.1; H,6.0; N,8.2; C₂₂H₂₇N₃O₅S,0.5C₂H₂O₄ requires : C,57.2; H,5.8; N,8.3%

Example 2

TABLE I

Compound	Formation of base				Salt formation				
	sulphonamide (g)	Indole (g)	Temp. (°C)	Time (h)	Yield (g)	Base (g)	Acid (g)	Solvent	Yield (g)
a	0.196	0.65	100-110	24	0.237	0.211	oxalic 0.065	EtOH	0.18
b	0.40	1.0	100	66	0.8	0.3	oxalic 0.09	EtOH	0.1
c	0.44	1.17	100	24	0.39	0.16	oxalic 0.045	EtOH	0.04
d	0.69	0.8	100	24	0.45	0.10	fumaric 0.015	EtOAc	0.056
e	0.65	0.273	120	17	0.43	0.153	oxalic 0.041	EtOH	0.147
f	0.44	1.17	100	24	0.66	0.10	fumaric 0.018	EtOAc	0.093
g	0.57	1.17	100	24	0.55	0.11	oxalic 0.027	EtOAc	0.075
h	0.74	0.74	110	16	0.38	0.097	oxalic 0.024	EtOH	0.025
i	0.25	0.39	100	24	0.16	0.16	oxalic 0.036	EtOAc	0.175
EtOH - Ethanol									
EtOAc - Ethyl acetate									

Example 3

4-[[[2-[3-[2-(Dimethylamino)ethyl]-1H-indol-5-yl]ethenyl]sulphonyl]amino]methyl]benzamide oxalate

A mixture of 5-iodo-N,N-dimethyl-1H-indole-3-ethanamine, oxalate (0.65g), 4-[[[ethenylsulphonyl]amino]methyl]benzamide (0.40g), palladium acetate (16mg) and triethylamine (0.7ml) in methanol (4ml) was heated in a 5ml "reacti-vial" at 100° for 22h. The mixture was evaporated to give an oil (1.65g) which was purified by column chromatography (B) to give a solid (245mg). This was dissolved in methanol (2ml) and a solution of oxalic acid (52mg) in methanol (2ml) was added. The mixture was evaporated to give a foam (288mg) which was recrystallised from ethanol/toluene and combined with similarly prepared material to afford the title compound as a solid, (312mg), m.p. 145-150°.

Analysis Found : C,56.4; H,5.4; N,9.7. C₂₄H₂₇N₃O₅S,0.10 EtOH, 0.32mol toluene requires C,57.65; H,5.8; N,10.2%.

Example 4

3-[2-(Dimethylamino)ethyl]-N-methyl-1H-indole-5-propanesulphonamide oxalate

A solution of the product of Example 1 (237.5mg) in absolute ethanol (20ml) was hydrogenated over pre-reduced 10% palladium oxide (450mg, 50% aqueous paste) at ambient temperature and pressure for a period of 24h. The reaction mixture was filtered through a celite-sand pad, which was washed thoroughly with ethanol (100ml) and the combined filtrates concentrated in vacuo. Flash chromatography (A) of the residue afforded the product as an oil (184.5mg), which was dissolved in absolute ethanol (1ml)

13

and filtered through a cotton wool plug. To this solution was added a solution of anhydrous oxalic acid (51.4mg) in absolute ethanol (0.50ml), and on scratching a crystalline material was deposited. The salt was filtered off, dried and recrystallised from absolute ethanol (5ml) to afford the *title compound* as an amorphous powder (80mg) m.p. 141-143° (softens 131°) Analysis Found; C,52.1; H,6.6; N,9.95.
 5 C₁₆H₁₇N₃O₃S.C₂H₄O₄ requires C,52.3; H,6.6; N,10.2%. N.m.r. δ (CD₂SOCD₂) 1.98(2H,m,CH₂CH₂SO₂NH) 2.53(d,MeNHSO₂), 2.83(6H,s, NMe₂), 2.7-3.35(8H,m,CH₂CH₂NMe₂ and CH₂CH₂SO₂NH), 6.85-7.45(5H,m, aromatic + NHSO₂).

Example 5

10 The following compounds were prepared according to the method of Example 4, using the starting materials and reaction conditions given in Table II below.

(a) 3-[2-(Dimethylamino)ethyl]-1H-indole-5-ethanesulphonamide oxalate
 m.p. 176-178° Analysis Found: C,49.45; H,5.9; N,10.8. C₁₈H₂₁N₃O₅S.C₂H₄O₄.0.32H₂O requires C,49.1; H,6.1; N,10.7%. N.m.r. δ (CD₂SOCD₂) 2.85(6H,s,NMe₂), 3.0-3.4(8H,m,CH₂CH₂SO₂NH, and CH₂CH₂NMe₂), 6.85-7.55(6H,m,aromatic + SO₂NH₂).

(b) 3-[2-(Dimethylamino)ethyl]-N-methyl-1H-indole-5-ethane-sulphonamide
 N.m.r. (CD₃OD) 2.42(6H,s,NMe₂), 2.74(5H,s,MeNHSO₂ and m, CH₂CH₂NMe₂), 2.98(2H,CH₂CH₂NMe₂), 3.16-3.44(4H,m,CH₂CH₂SO₂NHMe), 7.0-7.5(4H,m,aromatic).

(c) 3-[2-(Dimethylamino)ethyl]-N,N-dimethyl-1H-indole-5-ethane-sulphonamide oxalate
 m.p. 130-135°. Analysis Found: C,51.4; H,6.8; N,9.8. C₁₈H₂₁N₃O₅S.C₂H₄O₄.0.26H₂O requires C,51.7; H,6.6; N,10.05%. N.m.r. δ (CD₂SOCD₂) 2.81(12H,s,Me₂NSO₂ and CH₂NMe₂), 3.0-3.4 (8H,m,Me₂NSO₂CH₂CH₂ and CH₂CH₂NMe₂), 7.0-7.55(4H,m,aromatic).

(d) 3-[2-(Dimethylamino)ethyl]-N-(2-phenylethyl)-1H-indole-5-ethanesulphonamide oxalate
 m.p. 155-156° Analysis Found: C,58.5; H,6.4; N,8.3. C₂₀H₂₃N₃O₅S.C₂H₄O₄.0.08H₂O requires C,58.7; H,6.4; N,8.6%. N.m.r. δ (CD₂SOCD₂) 2.82(6H,s,NMe₂), 2.75-3.35(12H,m,-CH₂CH₂NMe₂ and -CH₂CH₂NHSO₂CH₂CH₂-), 6.95-7.5(10H,m, aromatic + NHSO₂).

(e) 3-[2-(Dimethylamino)ethyl]-N-(1-methylethyl)-1H-indole-5-ethane-sulphonamide oxalate
 m.p. 168-170° Analysis Found: C,53.3; H,6.8; N,9.8. C₁₉H₂₃N₃O₅S.C₂H₄O₄.0.1H₂O requires C,53.2; H,6.8; N,9.8%. N.m.r. δ (CD₂SOCD₂) 1.16(6H,d,CHMe), 2.82(6H,s, NMe₂), 3.0-3.35(8H,m,CH₂CH₂ NMe₂ and NHSO₂CH₂CH₂), 6.98-7.5(5H,aromatic + NHSO₂).

(f) 3-[2-(Dimethylamino)ethyl]-N-indole-5-ethane-sulphonamide oxalate
 m.p. 158-159°. Analysis Found: C,52.1; H,6.5; N,10.5. C₁₇H₁₉N₃O₅S.C₂H₄O₄.0.03H₂O requires C,52.2; H,6.6; N,10.1%. N.m.r. δ (CD₂SOCD₂) 1.12(3H,t,MeCH₂NHSO₂), 2.95-3.35(10H,m, MeCH₂NHSO₂CH₂CH₂ and CH₂CH₂NMe₂), 7.0-7.55(5H,m,aromatic + NHSO₂).

(g) N-Cyclopentyl-3-[2-(dimethylamino)ethyl]-1H-indole-5-ethanesulphonamide oxalate
 m.p. 181-182°. Analysis Found: C,55.4; H,7.0; N,8.9. C₂₀H₂₅N₃O₅S.C₂H₄O₄ requires C,55.5; H,6.9; N,9.2%. N.m.r. δ (CD₂SOCD₂) 1.4-1.96(8H,m,cyclopentyl CH₂ x 4) 2.83(6H,s,NMe₂), 3.0-3.36(8H,m,SO₂CH₂CH₂ and CH₂CH₂NMe₂), 3.65(1H,m,SO₂NHCH) 6.98-7.52(4H,m,aromatics).

(h) 3-[2-(Dimethylamino)ethyl]-N-[4-methoxyphenyl] methyl]-1H-indole-5-ethanesulphonamide oxalate
 m.p. 142-144° Analysis Found: C,55.9; H,6.2; N,8.0. C₂₁H₂₅N₃O₅S.C₂H₄O₄.0.5H₂O requires: C,56.0; H,6.3; N,8.2%. N.m.r. δ (CD₂SOCD₂) 2.83(6H,s,NMe₂), 2.9-3.35 (8H,m,CH₂CH₂SO₂NH and CH₂CH₂NMe₂), 3.75(3H,s,OMe), 4.15(2H,d,CH₂NHSO₂), 6.8-7.45(8H,m, aromatic).

Example 5

TABLE II

5	Starting material	Hydrogenation			Salt Formation				
Compound	Product of Ex.No.	Weight (g)	PdO/C (g)	Time (h)	Yield of base (g)	oxalic acid (g)	Solvent	Yield (g)	10
10									
a	2a	0.14	0.28	14	0.085	0.026	EtOH	0.053	
b	2b	0.056	0.11	6	0.045	-	-	-	
c	2c	0.25	0.95	18	0.22	0.010	EtOAc	0.040	
15 d	2d	0.31	0.60	18	0.137	0.031	(1) EtOAc (2) EtOH	0.040	15
e	2e	0.755	2.0	19	0.255	0.068	EtOH	0.274	
f	2f	0.56	0.9	18	0.20	0.030 (x2)	EtOH + MeOH	0.221	
g	2g	0.40	0.8	6h	0.234	0.051	EtOH	0.080	
20 h	2h	0.59	0.6	-	0.45	0.10	(1) EtOAc + MeOH + E (2) EtOH	0.25	20

25 EtOH - Ethanol
EtOAc - Ethyl acetate
MeOH - Methanol
E - Diethyl ether

30 Example 6

3-[2-(Dimethylamino)ethyl]-N-phenyl-1H-indole-5-ethanesulphonamide hemioxalate

A solution of the product of Example 2h (283mg) in absolute alcohol (30ml) was hydrogenated over pre-reduced 10% palladium oxide on charcoal (740mg, 50% aqueous paste pre-reduced in ethanol, 20ml) for a period of 18h at room temperature and pressure. The mixture was filtered through a sand-celite pad, which was thoroughly washed with ethanol (150ml). The filtrate was concentrated *in vacuo* and the residue was taken up in ethanol (30ml) and treated with Raney nickel (~50mg) for a period of 30min. The Raney nickel was removed by filtration, and the filtrate re-hydrogenated for a further 18h. The catalyst was removed by filtration through a sand-celite pad and the filtrate concentrated *in vacuo*. Flash chromatography (A) of the residue afforded the product as a low melting solid (103mg). A filtered solution of the solid in warm absolute ethanol (2ml) was added to an ethanolic solution of anhydrous oxalic acid (25mg in 1ml). On scratching, an amorphous solid was deposited, which was filtered off, air dried (1h) and recrystallized from ethanol (30ml) to afford the *title compound* as an amorphous powder, m.p. 144-146°.

Analysis Found: C, 58.9; H, 6.6; N, 9.0. $C_{20}H_{22}N_2O_5S$, 0.5C₂H₄O₄, 0.4C₂H₄O₄, 0.07H₂O requires C, 58.5; H, 6.7; N, 9.4%. N.m.r. δ (CD₃SOCD₃) 2.63(6H, s, NMe₂), 2.98(4H, m, CH₂CH₂NMe₂), 3.06(2H, m, SO₂CH₂), 3.38(2H, m, SO₂CH₂CH₂), 6.8-7.5(9H, m, aromatics).

The following examples illustrate pharmaceutical formulations according to the invention, containing (E)-2-[3-[2-(dimethylamino)ethyl]-1H-indol-5-yl]-N-[(4-methoxyphenyl)methyl]ethanesulphonamide may be formulated in a very similar manner.

50 Tablets for oral administration

Direct compression

		mg/tablet	
55	Active Ingredient	2.4	55
	Calcium hydrogen phosphate B.P.*	95.10	
	Croscarmellose sodium USP	2.00	
	Magnesium stearate, B.P.	0.50	
60	Compression weight	100mg	60

* of a grade suitable for direct compression

The active ingredient is sieved before use. The calcium hydrogen phosphate, croscarmellose sodium and active ingredient are weighed into a clean polythene bag. The powders are mixed by vigorous shaking then the magnesium stearate is weighed and added to the mix which is blended further. The mix is then compressed using a Manesty F3 tablet machine fitted with 5.5mm flat edge punches, into tablets 5 with target compression weight of 100mg.

Tablets may also be prepared by other conventional methods such as wet granulation.

Tablets of other strengths may be prepared by altering the ratio of active ingredient to lactose or the compression weight and using punches to suit.

The tablets may be film coated with suitable film forming materials, such as hydroxypropyl methylcellulose, using standard techniques. Alternatively the tablets may be sugar coated.

Capsules

		mg/capsule	
15	Active Ingredient	2.4	15
	*Starch 1500	196.6	
	Magnesium Stearate BP	1.00	
	Fill Weight	200.00	
20			20

* A form of directly compressible starch.

The active ingredient is sieved and blended with the excipients. The mix is filled into size No.2 hard gelatin capsules using suitable machinery. Other doses may be prepared by altering the fill weight and if necessary changing the capsule size to suit.

Syrup

		mg/5ml dose	
30	Active Ingredient	2.4	30
	Buffer)	
	Flavour)	
	Colour)	
	Preservative)	
35	Thickening agent)	35
	Sweetening agent)	
	Purified Water	to 5.00ml	

The active ingredient, buffer, flavour, colour, preservative, thickening agent and sweetening agent are dissolved in some water, the solution is adjusted to volume and mixed. The syrup produced is clarified by filtration.

45 Suppository for rectal administration

	Active Ingredient	2.4mg	
	*Witepsol H15	to 1.0g	
	*A proprietary grade of Adeps Solidus Ph. Eur.		
50			50

A suspension of the active ingredient in molten Witepsol is prepared and filled, using suitable machinery, into 1g size suppository moulds.

55 Injection for intravenous administration

		mg/ml	
	Active ingredient	0.6mg	
	Sodium Chloride BP	as required	
60	Water for Injection BP	to 1.0ml	60

Sodium chloride may be added to adjust the tonicity of the solution and the pH may be adjusted, using acid or alkali, to that of optimum stability and/or to facilitate solution of the active ingredient. Alternatively


tively suitable buffer salts may be used.

The solution is prepared, clarified and filled into appropriate size ampoules sealed by fusion of the glass. The injection is sterilised by heating in an autoclave using one of the acceptable cycles. Alternatively the solution may be sterilised by filtration and filled into sterile ampoules under aseptic conditions.

5 The solution may be packed under an inert atmosphere of nitrogen or other suitable gas.

CLAIMS

1. Indoles of the general formula (I):



wherein

R₁ represents a hydrogen atom or a C₁₋₆alkyl or C₃₋₆alkenyl group;


R₂ represents a hydrogen atom, a C₁₋₆alkyl, C₃₋₆alkenyl, or C₆₋₆ cycloalkyl group, or a phenyl or phenyl (C₁₋₆)alkyl group in which the phenyl ring may be unsubstituted or substituted by a halogen atom, a C₁₋₆alkyl, C₁₋₆alkoxy or hydroxyl group, or by a group -NR₅R₆, or -CONR₅R₆, wherein R₅ and R₆, which may be the same or different, each represents a hydrogen atom or a C₁₋₆alkyl or C₃₋₆alkenyl group, or together with the nitrogen atom to which they are attached form a saturated monocyclic 5 to 7-membered ring, which may contain an additional hetero function;

R₃ and R₄, which may be the same or different, each represents a hydrogen atom or a C₁₋₆alkyl or propenyl group or R₃ and R₄ together form an aralkylidene group;

Alk represents an alkyl chain containing two or three carbon atoms which may be unsubstituted or substituted by not more than two C₁₋₆alkyl groups; and


A' represents an alkenyl chain containing two to five carbon atoms, and salts and solvates thereof.

2. Indoles according to claim 1, wherein A' represents a group



wherein m is zero or an integer from 1 to 3 and n is zero or an integer from 1 to 3 and the sum of m and n does not exceed 3.

3. Indoles according to claim 1, represented by the general formula (I')




wherein R₁, R₂, R₃, R₄ and Alk are as defined for general formula (I) and m is zero or an integer from 1 to 3, and physiologically acceptable salts and solvates thereof.

4. Indoles according to any of claims 1 to 3, wherein Alk represents an unsubstituted alkyl chain containing two carbon atoms.

5. Indoles according to any of claims 1 to 4 in the E-configuration with regard to the double bond in the 5-substituent.

6. Indoles according to claim 1, of the general formula (Ia):



wherein

R_{1n} represents a hydrogen atom or a C₁₋₆alkyl group;

R_{2n} represents a hydrogen atom, a C₁₋₆alkyl group or a phenyl or phenyl (C₁₋₆)alkyl group in which the phenyl ring is unsubstituted or substituted by a C₁₋₆alkoxy group or by the group -CONH₁;

R_{3n} and R_{4n} each represents a hydrogen atom or a C₁₋₆alkyl group; and

ma is zero or 1;


and physiologically acceptable salts and solvates thereof.

7. Indoles according to claim 1, selected from
 (E)-2-[3-[2-(dimethylamino)ethyl]-1H-indol-5-yl]-N-methylethanesulphonamide;
 (E)-2-[3-[2-(dimethylamino)ethyl]-1H-indol-5-yl]-N-(2-phenylethyl)ethanesulphonamide;
 (E)-2-[3-[2-(dimethylamino)ethyl]-1H-indol-5-yl]-N-[(4-methoxyphenyl)methyl]ethanesulphonamide;

5 and the physiologically acceptable salts and solvates thereof.

8. A pharmaceutical composition which comprises as active ingredient an effective amount of at least one indole of general formula (I) according to claim 1 or a physiologically acceptable salt or solvate thereof together with one or more pharmaceutically acceptable carriers or excipients.

9. A process for the preparation of a compound of general formula (II):




wherein R₁, R₂, R₃, R₄ and Alk are as defined in claim 1, and A represents an alkyl chain containing two to five carbon atoms which comprises reducing an indole of general formula (I) as defined in claim 1.

10. Compounds of general formula (II) as defined in claim 9, selected from


- 20 3-[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-propanesulphonamide;
 3-[2-(dimethylamino)ethyl]-N,N-dimethyl-1H-indole-5-ethanesulphonamide;
 3-[2-(dimethylamino)ethyl]-N-(2-phenylethyl)-1H-indole-5-ethanesulphonamide;
 3-[2-(dimethylamino)ethyl]-N-(1-methylethyl)-1H-indole-5-ethanesulphonamide;
 3-[2-(dimethylamino)ethyl]-N-ethyl-1H-indole-5-ethanesulphonamide;
 25 3-[2-(dimethylamino)ethyl]-N-phenyl-1H-indole-5-ethanesulphonamide; and
 N-cyclopentyl-3-[2-(dimethylamino)ethyl]-1H-indole-5-ethanesulphonamide.

11. A process for the preparation of an indole of general formula (I) according to claim 1 or a salt or solvate thereof which comprises:


(A) reacting an indole of general formula (III):




wherein X represents a leaving atom or group and Alk, R₃ and R₄ are as defined in claim 1 with an alkene of formula (IV):




wherein -A'-CH₂ represents a C₁₋₅ alkenyl chain and R₁ and R₂ are as defined in claim 1; or
 (B) reacting an aldehyde of formula (V):




wherein A' represents a direct bond or a C₁₋₅ alkyl chain and Alk, R₃ and R₄ are as defined in claim 1, with a reagent serving to form the group R₁R₂NSO₂A' wherein R₁, R₂ and A' are as defined in claim 1; or
 (C) subjecting a compound of general formula (X):




wherein R₁, R₂, R₃, R₄ and Alk are as defined in claim 1 and A⁶ represents a C₁₋₅ alkyl chain substituted by a leaving atom or group, X¹, to a reaction to eliminate HX¹, or
 (D) cyclising a compound of general formula (XI):



5 wherein R_1 , R_2 , Alk and A^1 are as defined in claim 1 and Q is the group NR_3R_4 (where R_3 and R_4 are as defined in claim 1) or a protected derivative thereof or a leaving atom or group; or
(E) reacting a compound of general formula (XIV):



15 wherein R_1 , R_2 , A^1 and Alk are as defined in claim 1 and Y is a readily displaceable atom or group or a protected derivative thereof with an amine of formula R_3R_4NH (where R_3 and R_4 are as defined in claim 1); or
(F) reacting a compound of general formula (XV):



25 wherein A^1 , Alk, R_3 and R_4 are as defined in claim 1 and Z is a leaving atom or group with a compound of general formula (XVI);



wherein R_1 and R_2 are as defined in claim 1; or

35 (G) converting a compound of general formula (I) as defined in claim 1 or a salt or protected derivative thereof into another compound of general formula (I); or

(H) subjecting a protected derivative of general formula (I) as defined in claim 1 or a salt thereof to reaction to remove the protecting group or groups; and if necessary and/or desired effecting one or two additional reactions subsequent to any of processes A' to G comprising:-

- 40
- (i) removing any protecting group or groups; and
 - (ii) converting a compound of general formula (I) or a salt thereof into a physiologically acceptable salt or solvate thereof.